

# תוכנית טיפול DYSPOORT

שם המטופל: \_\_\_\_\_ ת.ז. \_\_\_\_\_

## CERVICAL DYSTONIA

### LEFT ANTEROLATERAL

מינון בפועל	שם השריר
	Sternocleidomastoid
	Scalenus medius*

### RIGHT ANTEROLATERAL

מינון בפועל	שם השריר
	Sternocleidomastoid
	Scalenus medius*

### LEFT UPPER POSTERIOR

מינון בפועל	שם השריר
	Trapezius pars descendens
	Splenius capitis
	Longissimus capitis*
	Semispinalis capitis*
	Obliquus capitis inferior*

### RIGHT UPPER POSTERIOR

מינון בפועל	שם השריר
	Trapezius pars descendens
	Splenius capitis
	Longissimus capitis*
	Semispinalis capitis*
	Obliquus capitis inferior*

### LEFT LOWER POSTERIOR

מינון בפועל	שם השריר
	Splenius cervicis
	Longissimus cervicis*
	Semispinalis cervicis*
	Trapezius transverse
	Levator scapulae


### RIGHT LOWER POSTERIOR

מינון בפועל	שם השריר
	Splenius cervicis
	Longissimus cervicis*
	Semispinalis cervicis*
	Trapezius transverse
	Levator scapulae

\*Muscle not in Dysport indication

#### Dosing guidance in Blepharospasm / Hemifacial Spasm

מינון בפועל	טווח מינון דיספורט	שם השריר
	10U	Upper medial orbicularis oculi muscle
	10U	Upper lateral orbicularis oculi muscle
	10U	Lower medial orbicularis oculi muscle
	10U	Lower lateral orbicularis oculi muscle
	10U	Frontalis muscles



A dose of 40 units per eye was significantly effective. A dose of 80 to maximum of 120 units per eye resulted in a longer duration of effect.


Resulting Dose Unit per ml	Diluent* per 500 Unit Vial	Diluent* per 300 Unit Vial
500 Units	1 ml	0.6 ml
200 Units	2.5 ml	1.5 ml
100 Units	5 ml	3 ml

\*Preservative-free 0.9% sodium chloride injection

#### Example of Reconstitution:<sup>1</sup>



500U Vial  
Dysport®



2.5mL preservative-free 0.9%  
Sodium Chloride



**Concentration:**  
20 Units  
per 0.1mL



The maximum dose administered in a single session involving both the upper and lower limbs must not exceed 1500 Units.<sup>1</sup>

The units of Dysport® (BOTULINUM A TOXIN) are specific and are not interchangeable with other products which contain botulinum toxin.

Dysport® (BOTULINUM A TOXIN) is available in 300U and 500U vial sizes. Vial size selected based on the maximum dose required by your patient.

**Dysport®** (aboBoNT-A) offers a duration of response of 12 - 16 weeks or more<sup>1</sup>

**Dysport®** (aboBoNT-A) gives your patients long-lasting symptom relief<sup>1</sup> and a chance of feeling a little more like themselves between injections

For full details please refer to the prescribing information as approved by the Israeli MoH

1. Botulinum A toxin (Dysport®) prescribing information as approved by the Israeli MoH (03.2021)